



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|--|--|---|---|
| (51) International Patent Classification ⁶ : C07D 209/42, 209/52, 401/12, 413/04, 405/12, 409/12, 471/04, A61K 31/44, 31/40 | | A1 | (11) International Publication Number: WO 95/11885 |
| | | | (43) International Publication Date: 4 May 1995 (04.05.95) |
| (21) International Application Number: PCT/US94/12300 | | (81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BI, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). | |
| (22) International Filing Date: 26 October 1994 (26.10.94) | | | |
| (30) Priority Data: 08/144,138 27 October 1993 (27.10.93) US | | | |
| (60) Parent Application or Grant (63) Related by Continuation US 08/144,138 (CIP) Filed on 27 October 1994 (27.10.94) | | Published With international search report. | |
| (71) Applicant (for all designated States except US): NEUROGEN CORPORATION (US/US); 35 N.E. Industrial Road, Bran- ford, CT 06405 (US). | | | |
| (72) Inventors; and | | | |
| (75) Inventors/Applicants (for US only): ALBAUGH, Pamela (US/US); 81 Long Hill Road, Clinton, CT 06413 (US); HUTCHISON, Alan (US/US); 175 Bartlett Drive, Madison, CT 06443 (US). | | | |
| (74) Agent: SARUSS, Steven, J.; Allegretti & Witcoff, Ltd., Ten South Wacker Drive, Chicago, IL 60606 (US). | | | |
| (54) Title: CERTAIN FUSED PYRROLECARBOXANILIDES; A NEW CLASS OF GABA BRAIN RECEPTOR LIGANDS | | | |
| | | | |
| (57) Abstract | | | |
| <p>The present invention encompasses structures of formula (I) and the pharmaceutically acceptable non-toxic salts thereof wherein (a) represents (b), (c), (d) or (e), wherein W represents substituted or unsubstituted aryl groups; X is hydrogen, hydroxy or lower alkyl; T is hydrogen, halogen, hydroxyl, amino or alkyl; R₁ is hydrogen or an organic group; R₄ is hydrogen or substituted or unsubstituted organic substituent; R₅ and R₆ represent organic, and inorganic substituents; and n is 1, 2, 3, or 4. These compounds are highly selective agonists, antagonists or inverse agonists for GABA_A brain receptors or prodrugs of agonists, antagonists or inverse agonists for GABA_A brain receptors. These compounds are useful in the diagnosis and treatment of anxiety, sleep and seizure disorders, overdose with benzodiazepine drugs and for enhancement of memory.</p> | | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|---------------------------------------|----|--------------------------|
| AT | Austria | GB | United Kingdom | MR | Mauritania |
| AU | Australia | GE | Georgia | MW | Malawi |
| BB | Barbados | GN | Guinea | NE | Niger |
| BE | Belgium | GR | Greece | NL | Netherlands |
| BF | Burkina Faso | HU | Hungary | NO | Norway |
| BG | Bulgaria | IE | Ireland | NZ | New Zealand |
| BJ | Benin | IT | Italy | PL | Poland |
| BR | Brazil | JP | Japan | PT | Portugal |
| BY | Belarus | KE | Kenya | RO | Romania |
| CA | Canada | KG | Kyrgyzstan | RU | Russian Federation |
| CF | Central African Republic | KP | Democratic People's Republic of Korea | SD | Sudan |
| CG | Congo | KR | Republic of Korea | SE | Sweden |
| CH | Switzerland | KZ | Kazakhstan | SI | Slovenia |
| CI | Côte d'Ivoire | LI | Liechtenstein | SK | Slovakia |
| CM | Cameroun | LK | Sri Lanka | SN | Senegal |
| CN | China | LU | Luxembourg | TD | Chad |
| CS | Czechoslovakia | LV | Latvia | TG | Togo |
| CZ | Czech Republic | MC | Monaco | TJ | Tajikistan |
| DE | Germany | MD | Republic of Moldova | TT | Trinidad and Tobago |
| DK | Denmark | MG | Madagascar | UA | Ukraine |
| ES | Spain | MN | Mongolia | US | United States of America |
| FI | Finland | | | UZ | Uzbekistan |
| FR | France | | | VN | Viet Nam |
| GA | Gabon | | | | |

5 **Certain Fused Pyrrolicarboxanilides; A New Class of
GABA Brain Receptor Ligands**

BACKGROUND OF THE INVENTION

10 This application is a continuation-in-part of U.S. Application Serial
No. 08/144,138, filed October 27, 1993.

Field of the Invention

15 This invention relates to certain fused pyrrolicarboxanilides which
selectively bind to GABA_A receptors. This invention also relates to
pharmaceutical compositions comprising such compounds. It further
relates to the use of such compounds in treating anxiety, sleep and seizure
disorders, and overdoses of benzodiazepine-type drugs, and enhancing
alertness. The interaction of fused pyrrolicarboxanilides of the invention
20 with a GABA binding site, the benzodiazepines (BDZ) receptor, is described.
This interaction results in the pharmacological activities of these
compounds.

Description of the Related Art

25 γ -Aminobutyric acid (GABA) is regarded as one of the major
inhibitory amino acid transmitters in the mammalian brain. Over 30 years
have elapsed since its presence in the brain was demonstrated (Roberts &
Frankel, J. Biol. Chem. 187: 55-63, 1950; Udenfriend, J. Biol. Chem. 187: 65-69,
1950). Since that time, an enormous amount of effort has been devoted to
30 implicating GABA in the etiology of seizure disorders, sleep, anxiety and
cognition (Tallman and Gallager, Ann. Rev. Neuroscience 8: 21-44, 1985).
Widely, although unequally, distributed through the mammalian brain,
GABA is said to be a transmitter at approximately 30% of the synapses in the
brain. In most regions of the brain, GABA is associated with local inhibitory
35 neurons and only in two regions is GABA associated with longer projections.
GABA mediates many of its actions through a complex of proteins localized
both on cell bodies and nerve endings; these are called GABA_A receptors.
Postsynaptic responses to GABA are mediated through alterations in chloride
conductance that generally, although not invariably, lead to
40 hyperpolarization of the cell. Recent investigations have indicated that the
complex of proteins associated with postsynaptic GABA responses is a major

site of action for a number of structurally unrelated compounds capable of modifying postsynaptic responses to GABA. Depending on the mode of interaction, these compounds are capable of producing a spectrum of activities (either sedative, anxiolytic, and anticonvulsant, or wakefulness, seizures, and anxiety).

1,4-Benzodiazepines continue to be among the most widely used drugs in the world. Principal among the benzodiazepines marketed are chlordiazepoxide, diazepam, flurazepam, and triazolam. These compounds are widely used as anxiolytics, sedative-hypnotics, muscle relaxants, and anticonvulsants. A number of these compounds are extremely potent drugs: such potency indicates a site of action with a high affinity and specificity for individual receptors. Early electrophysiological studies indicated that a major action of benzodiazepines was enhancement of GABAergic inhibition. The benzodiazepines were capable of enhancing presynaptic inhibition of a monosynaptic ventral root reflex, a GABA-mediated event (Schmidt et al., 1967, Arch. Exp. Path. Pharmacol. 258: 69-82). All subsequent electrophysiological studies (reviewed in Tallman et al. 1980, Science 207: 274-81, Haefley et al., 1981, Handb. Exptl. Pharmacol. 33: 95-102) have generally confirmed this finding, and by the mid-1970s, there was a general consensus among electrophysiologists that the benzodiazepines could enhance the actions of GABA.

With the discovery of the "receptor" for the benzodiazepines and the subsequent definition of the nature of the interaction between GABA and the benzodiazepines, it appears that the behaviorally important interactions of the benzodiazepines with different neurotransmitter systems are due in a large part to the enhanced ability of GABA itself to modify these systems. Each modified system, in turn, may be associated with the expression of a behavior.

Studies on the mechanistic nature of these interactions depended on the demonstration of a high-affinity benzodiazepine binding site (receptor). Such a receptor is present in the CNS of all vertebrates phylogenetically newer than the bony fishes (Squires & Braestrup 1977, Nature 166: 732-34, Mohler & Okada, 1977, Science 198: 854-51, Mohler & Okada, 1977, Br. J. Psychiatry 133: 261-68). By using tritiated diazepam, and a variety of other compounds, it has been demonstrated that these benzodiazepine binding sites fulfill many of the criteria of pharmacological receptors; binding to these sites *in vitro* is rapid, reversible, stereospecific, and saturable. More importantly, highly significant correlations have been shown between the ability of benzodiazepines to displace diazepam from its binding site and activity in a number of animal behavioral tests predictive of benzodiazepine

potency (Braestrup & Squires 1978, Br. J. Psychiatry 133: 249-60, Mohler & Okada, 1977, Science 198: 854-51, Mohler & Okada, 1977, Br. J. Psychiatry 133: 261-68). The average therapeutic doses of these drugs in man also correlate with receptor potency (Tallman et al. 1980, Science 207: 274-281).

- 5 In 1978, it became clear that GABA and related analogs could interact at the low affinity (1 mM) GABA binding site to enhance the binding of benzodiazepines to the clonazepam-sensitive site (Tallman et al. 1978, Nature, 274: 383-85). This enhancement was caused by an increase in the affinity of the benzodiazepine binding site due to occupancy of the GABA site. The data
10 were interpreted to mean that both GABA and benzodiazepine sites were allosterically linked in the membrane as part of a complex of proteins. For a number of GABA analogs, the ability to enhance diazepam binding by 50% of maximum and the ability to inhibit the binding of GABA to brain membranes by 50% could be directly correlated. Enhancement of benzodiazepine
15 binding by GABA agonists is blocked by the GABA receptor antagonist (+) bicuculline; the stereoisomer (-) bicuculline is much less active (Tallman et al., 1978, Nature, 274: 383-85).

- Soon after the discovery of high affinity binding sites for the benzodiazepines, it was discovered that a triazolopyridazine could interact
20 with benzodiazepine receptors in a number of regions of the brain in a manner consistent with receptor heterogeneity or negative cooperativity. In these studies, Hill coefficients significantly less than one were observed in a number of brain regions, including cortex, hippocampus, and striatum. In cerebellum, triazolopyridazine interacted with benzodiazepine sites with
25 a Hill coefficient of 1 (Squires et al., 1979, Pharma. Biochem. Behav. 10: 825-30, Klepner et al. 1979, Pharmacol. Biochem. Behav. 11: 457-62). Thus, multiple benzodiazepine receptors were predicted in the cortex, hippocampus, striatum, but not in the cerebellum.

- Based on these studies, extensive receptor autoradiographic
30 localization studies were carried out at a light microscopic level. Although receptor heterogeneity has been demonstrated (Young & Kuhar 1980, J. Pharmacol. Exp. Ther. 212: 337-46, Young et al., 1981 J. Pharmacol. Exp. Ther. 216: 425-430, Niehoff et al. 1982, J. Pharmacol. Exp. Ther. 221: 670-75), no simple correlation between localization of receptor subtypes and the
35 behaviors associated with the region has emerged from the early studies. In addition, in the cerebellum, where one receptor was predicted from binding studies, autoradiography revealed heterogeneity of receptors (Niehoff et al., 1982, J. Pharmacol. Exp. Ther. 221: 670-75).

- A physical basis for the differences in drug specificity for the two
40 apparent subtypes of benzodiazepine sites has been demonstrated by

Seighart & Karobath, 1980, *Nature* 286: 285-87. Using gel electrophoresis in the presence of sodium dodecyl sulfate, the presence of several molecular weight receptors for the benzodiazepines has been reported. The receptors were identified by the covalent incorporation of radioactive flunitrazepam, a benzodiazepine which can covalently label all receptor types. The major labeled bands have molecular weights of 50,000 to 53,000, 55,000, and 57,000 and the triazolopyridazines inhibit labeling of the slightly higher molecular weight forms (53,000, 55,000, 57,000) (Seighart et al. 1983, *Eur. J. Pharmacol.* 88: 291-99).

At that time, the possibility was raised that the multiple forms of the receptor represent "isoreceptors" or multiple allelic forms of the receptor (Tallman & Gallager 1985, *Ann. Rev. Neurosci.* 8, 21-44). Although common for enzymes, genetically distinct forms of receptors have not generally been described. As we begin to study receptors using specific radioactive probes and electrophoretic techniques, it is almost certain that isoreceptors will emerge as important in investigations of the etiology of psychiatric disorders in people.

The GABA_A receptor subunits have been cloned from bovine and human cDNA libraries (Schoenfield et al., 1988; Duman et al., 1989). A number of distinct cDNAs were identified as subunits of the GABA_A receptor complex by cloning and expression. These are categorized into α , β , γ , δ , ϵ , and provide a molecular basis for the GABA_A receptor heterogeneity and distinctive regional pharmacology (Shivvers et al., 1980; Levitan et al., 1989). The γ subunit appears to enable drugs like benzodiazepines to modify the GABA responses (Pritchett et al., 1989). The presence of low Hill coefficients in the binding of ligands to the GABA_A receptor indicates unique profiles of subtype specific pharmacological action.

Drugs that interact at the GABA_A receptor can possess a spectrum of pharmacological activities depending on their abilities to modify the actions of GABA. For example, the beta-carbolines were first isolated based upon their ability to inhibit competitively the binding of diazepam to its binding site (Nielsen et al., 1979, *Life Sci.* 25: 679-86). The receptor binding assay is not totally predictive about the biological activity of such compounds; agonists, partial agonists, inverse agonists, and antagonists can inhibit binding. When the beta-carboline structure was determined, it was possible to synthesize a number of analogs and test these compounds behaviorally. It was immediately realized that the beta-carbolines could antagonize the actions of diazepam behaviorally (Tenen & Hirsch, 1980, *Nature* 288: 609-10). In addition to this antagonism, beta-carbolines possess intrinsic activity of


their own opposite to that of the benzodiazepines: they become known as inverse agonists.

In addition, a number of other specific antagonists of the benzodiazepine receptor were developed based on their ability to inhibit the binding of benzodiazepines. The best studied of these compounds is an imidazodiazepine (Hunkeler et al., 1981, Nature 290: 514-516). This compound is a high affinity competitive inhibitor of benzodiazepine and beta-carboline binding and is capable of blocking the pharmacological actions of both these classes of compounds. By itself, it possesses little intrinsic pharmacological activity in animals and humans (Hunkeler et al., 1981, Nature 290: 514-16; Darragh et al., 1983, Eur. J. Clin. Pharmacol. 14: 569-70). When a radiolabeled form of this compound was studied (Mohler & Richards, 1981, Nature 294: 763-65), it was demonstrated that this compound would interact with the same number of sites as the benzodiazepines and beta-carbolines, and that the interactions of these compounds were purely competitive. This compound is the ligand of choice for binding to GABA_A receptors because it does not possess receptor subtype specificity and measures each state of the receptor.


The study of the interactions of a wide variety of compounds similar to the above has led to the categorizing of these compounds. Presently, those compounds possessing activity similar to the benzodiazepines are called agonists. Compounds possessing activity opposite to benzodiazepines are called inverse agonists, and the compounds blocking both types of activity have been termed antagonists. This categorization has been developed to emphasize the fact that a wide variety of compounds can produce a spectrum of pharmacological effects, to indicate that compounds can interact at the same receptor to produce opposite effects, and to indicate that beta-carbolines and antagonists with intrinsic anxiogenic effects are not synonymous.

A biochemical test for the pharmacological and behavioral properties of compounds that interact with the benzodiazepine receptor continues to emphasize the interaction with the GABAergic system. In contrast to the benzodiazepines, which show an increase in their affinity due to GABA (Tallman et al., 1978, Nature 274: 383-85, Tallman et al., 1980, Science 207: 274-81), compounds with antagonist properties show little GABA shift (i.e., change in receptor affinity due to GABA) (Mohler & Richards 1981, Nature 294: 763-65), and the inverse agonists actually show a decrease in affinity due to GABA (Braestrup & Nielson 1981, Nature 294: 472-474). Thus, the GABA shift predicts generally the expected behavioral properties of the compounds.

Various compounds have been prepared as benzodiazepine agonists and antagonists. For Example, U.S. Patents Nos. 3,455,943, 4,435,403, 4,596,808, 4,623,649, and 4,719,210, German Patent No. DE 3,246,932, and Liebig's Ann. Chem. 1986, 1749 teach assorted benzodiazepine agonists and antagonists and related anti-depressant and central nervous system active compounds. U.S. Patent No. 3,455,943 discloses compounds of the formula:



wherein R_1 is a member of the group consisting of hydrogen and lower alkoxy; R_2 is a member of the group consisting of hydrogen and lower alkoxy; R_3 is a member of the group consisting of hydrogen and lower alkyl; and X is a divalent radical selected from the group consisting of




and



and the non-toxic acid addition salts thereof.

U.S. Patent No. 4,435,403 teaches compounds of the formula:




wherein

R^C is hydrogen, lower alkyl, alkoxyalkyl of up to 6 C-atoms, cycloalkyl of 3-6 C-atoms, arylalkyl of up to 8 C-atoms, or $(CH_2)_nOR_2$ wherein R_{20} is alkyl of up to 6 C-atoms, cycloalkyl of 3-6 C-atoms or arylalkyl of up to


8 C-atoms and n is an integer of 1 to 3; Y is oxygen, two hydrogen atoms or NOR₁, wherein R₁ is hydrogen, lower alkyl, aryl or arylalkyl of up to 6 C-atoms, COR₂, wherein R₂ is lower alkyl of up to 6 C-atoms, or Y is CHCOOR₃, wherein R₃ is hydrogen or lower alkyl or Y is NNR₄R₅, wherein R₄ and R₅ can be the same or different and each is hydrogen, lower alkyl, C₆-10-aryl, C₇-10-arylalkyl or CONR₆R₇, wherein R₆ and R₇ can be the same or different and each is hydrogen or lower alkyl or R₄ and R₅ together with the connecting N-atom, form a 5- or 6-membered heterocyclic ring which optionally may also contain an O-atom or up to 3 N-atoms and which optionally may be substituted by a lower alkyl group; Z is hydrogen, or alkoxy or aralkoxy each of up to 10 C-atoms and each optionally substituted by hydroxy, or Z is alkyl of up to 6 C-atoms, C₆-10-aryl or C₇-10-arylalkyl each of which may optionally be substituted by a COOR₈ or a CONR₉R₁₀ group, wherein R₈ is alkyl of up to 6 C-atoms, and R₉ and R₁₀ can be the same or different and each is hydrogen or alkyl of up to 6 C-atoms; or Z is NR₉R₁₀, wherein R₉ and R₁₀ are as defined above; or Z is NR₁₁CHR₁₂R₁₃, wherein R₁₁ and R₁₂ each is hydrogen or together form a N=C double bond, wherein R₁₃ is C₁-10-alkyl or NR₁₄R₁₅, wherein R₁₄ and R₁₅ are the same or different and each is hydrogen, OH or alkyl or alkoxy each of up to 6 C-atoms, or wherein R₁₂ and R₁₃ together are oxygen, in which case, R₁₁ is hydrogen; or Z is COOR₂ wherein R₂ is as defined above; or Y and Z, together with the connecting C-atom, may form a 5- or 6-membered heterocyclic ring which contains an O-atom, adjoining O- and N-atoms or up to 4 N atoms and which optionally may be substituted by a lower alkyl group, hydroxy or oxo.

U.S. Patent No. 4,596,808 discloses compounds of the formula:




wherein R^A is H, F, Cl, Br, I, NO_2 , CN, CH_3 , CF_3 , SCH_3 , $\text{NR}_{16}\text{R}_{17}$ or NHCOR_{16} , wherein R_{16} of R_{17} are the same or different and each is hydrogen or alkyl, alkenyl or alkynyl each of up to 6 C-atoms, arylalkyl or cycloalkyl each of up to 10 C-atoms, or wherein R_{16} and R_{17} together form a saturated or unsaturated 3-7 membered heterocyclic ring.

U.S. Patent No. 4,623,649 teaches compounds of the formula:




wherein R_3 is an oxadiazolyl residue of the formula



wherein R_5 stands for lower alkyl of up to 3 carbon atoms or an ester $-\text{CO}_2\text{R}_6$ with R_6 being hydrogen or lower alkyl of up to 3 carbon atoms. R_4 is hydrogen, lower alkyl of up to 3 carbon atoms, or CH_2OR_9 wherein R_9 is lower alkyl of up to 3 carbon atoms, R^A is phenyl or a hydrocarbon residue containing 2-10 carbon atoms which can be cyclic or acyclic, saturated or unsaturated, branched or unbranched, and which can optionally be substituted by oxo, formyl OH, O-alkyl of up to 3 carbon atoms or phenyl, and wherein in a cyclic hydrocarbon residue, a CH_2 -group can be replaced by oxygen.


U.S. Patent No. 4,719,210 discloses compounds of the formula:



wherein R_1 is hydrogen or a protecting group, R_2 is $-CH=CR_4$ or $-C=CR_4$, R_4 is hydrogen or halogen, R_3 is hydrogen, lower alkyl or lower alkoxyalkyl, R^A is, *inter alia*, hydrogen, OR₇, lower alkyl, which optionally is substituted with aryl, lower alkoxy or NR₅R₆, R_5 and R_6 can be the same or different and in each case is hydrogen, lower alkyl or together with the nitrogen atom a 5-6 member ring, which can contain another heteroatom. R_7 is lower alkyl, optionally substituted aryl or arylalkyl, and each compound can contain one or more R^A radicals which are not hydrogen.

These compounds differ from the compounds of the present invention. These U.S. Patents teach carbocyclic compounds having pyridine or piperidine rings but lacking the pyrimidine ring present in the compounds of the present invention.

German Patent No. DE 3,246,932 discloses compounds of the formula:



20


wherein

$R =$ halogen, NO_2 , CO_2H , modified CO_2H , R_2O , $R_2S(O)_n$; $n = 0-2$; and $R_1 = H$, alkyl, cycloalkyl, arylalkyl, aryl, CO_2H , amino R_2O , $R_2S(O)_n$. However no examples were exemplified in this patent with $R_1=aryl$.

25

Liebigs Ann. Chem. 1986. 1749-1764 teaches compounds of the formula:

30







where R^X is hydrogen, methyl, benzyloxy, or methoxy, and R_3 is carboethoxy.


- 5 None of these compounds are indole-3-carboxamides and no such compounds displaying activity at GABA receptors have been described.

A variety of indole-3-carboxamides are described in the literature. For example, J. Org. Chem., 42: 1883-1885 (1977) discloses the following compounds.

10



J. Heterocyclic Chem., 14: 519-520 (1977) discloses a compound of the following formula:




- 5 None of these indole-3-carboxamides includes an oxy substituent at the 4-position of the indole ring.

12
SUMMARY OF THE INVENTION

This invention provides novel compounds of Formula I which interact with a GABAA binding site, the benzodiazepine receptor.

The invention provides pharmaceutical compositions comprising compounds of Formula I. The invention also provides compounds useful in the diagnosis and treatment of anxiety, sleep and seizure disorders, overdose with benzodiazepine drugs and for enhancement of memory. Accordingly, a broad embodiment of the invention is directed to compounds of general Formula I:



I

15

or the pharmaceutically acceptable non-toxic salts thereof wherein:

T is

20

halogen, hydrogen, hydroxyl, amino or straight or branched chain lower alkoxy having 1-6 carbon atoms;

X is

25



hydrogen, hydroxyl or straight or branched chain lower alkyl having 1-6 carbon atoms;

W is

30

phenyl, 2 or 3-thienyl, 2, 3, or 4-pyridyl or 6-quinoliny, each of which may be mono or disubstituted with halogen, cyano, hydroxy, straight or branched chain lower alkyl having 1-6 carbon atoms, amino, mono or dialkylamino where each alkyl is independently straight or branched chain lower alkyl having 1-6 carbon atoms, straight or branched chain lower alkoxy having 1-6 carbon atoms, or NR_1COR_2 , COR_2 , CONR_1R_2 or CO_2R_2 where R_1 and R_2 are the same or different and represent hydrogen or straight or branched chain lower alkyl having 1-6 carbon atoms; and

35



5

wherein:

Y represents nitrogen or C-R₄;

10

Z represents N-R₇ or a carbon atom substituted with R₈ and R₉, i.e., C(R₈)(R₉);

n is 1, 2, 3, or 4;

15

R₃ is

hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2, 3 or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms;

20

R₄ is

halogen or trifluoromethyl; or

25

-OR₁₀, -COR₁₀, -CO₂R₁₀, -OCOR₁₀, or -R₁₀, where R₁₀ is hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2, 3 or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; or

30

-CONR₁₁R₁₂ or -(CH₂)_mNR₁₁R₁₂, where m is 0, 1, or 2; R₁₁ represents hydrogen, straight or branched chain lower alkyl having 1-6 carbon atoms; and R₁₂ is hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or branched chain lower alkyl

14
having 1-6 carbon atoms, or phenylalkyl or 2, 3 or 4-
pyridylalkyl where each alkyl is straight or branched chain
lower alkyl having 1-6 carbon atoms; or NR₁₁R₁₂ forms a
heterocyclic group which is morpholyl, piperidyl, pyrrolidyl,
or N-alkyl piperazyl;

R₅ and R₆ are the same or different and represent

hydrogen, halogen, straight or branched chain lower
alkyl having 1-6 carbon atoms, or straight or branched chain
lower alkoxy having 1-6 carbon atoms;

R₇ is

hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or
branched chain lower alkyl having 1-6 carbon atoms, or
phenylalkyl or 2,3, or 4-pyridylalkyl where each alkyl is
straight or branched chain lower alkyl having 1-6 carbon
atoms;

R₈ is

hydrogen or straight or branched chain lower alkyl
having 1-6 carbon atoms; and

R₉ is

-COR₁₃, -CO₂R₁₃ or -R₁₃, where R₁₃ is hydrogen, phenyl,
2, 3 or 4-pyridyl, straight or branched chain lower alkyl
having 1-6 carbon atoms, or phenylalkyl or 2, 3, or 4-
pyridylalkyl where each alkyl is straight or branched chain
lower alkyl having 1-6 carbon atoms; or

-CONR₁₄R₁₅ or -(CH₂)_kNR₁₄R₁₅, where k is 0, 1, or 2;
R₁₄ represents hydrogen, straight or branched chain lower
alkyl having 1-6 carbon atoms; and R₁₅ is hydrogen, phenyl,
2, 3, or 4-pyridyl, straight or branched chain lower alkyl
having 1-6 carbon atoms, or phenylalkyl or 2, 3 or 4-
pyridylalkyl where each alkyl is straight or branched chain
lower alkyl having 1-6 carbon atoms; or NR₁₄R₁₅ forms a
heterocyclic group which is morpholyl, piperidyl, pyrrolidyl,
or N-alkyl piperazyl.

These compounds are highly selective agonists, antagonists or inverse agonists for GABA_A brain receptors or prodrugs of agonists, antagonists or inverse agonists for GABA_A brain receptors. In other words, while the compounds of the invention, all interact with GABA_A brain receptors, they do display identical physiologic activity. Thus, these compounds are useful in the diagnosis and treatment of anxiety, sleep and seizure disorders, overdose with benzodiazepine drugs and for enhancement of memory. For example, these compounds can easily be used to treat overdoses of benzodiazepine-type drugs as they would competitively bind to the benzodiazepine receptor.


16
BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1-20 show representative fused pyrrolicarboxanilides of the present invention.

17
DETAILED DESCRIPTION OF THE INVENTION

The novel compounds encompassed by the instant invention can be described by general formula I set forth above or the pharmaceutically acceptable non-toxic salts thereof.


In addition, the present invention encompasses compounds of Formula II.




II

or the pharmaceutically acceptable non-toxic salts thereof wherein W, R₈ and R₉ are as defined above.

The present invention also encompasses compounds of Formula IIIa and IIIb:




IIIa



IIIb

or the pharmaceutically acceptable non-toxic salts thereof wherein W, R₃, and R₅ are as defined above.

The present invention also encompasses compounds of Formula IV:




IV

or the pharmaceutically acceptable non-toxic salts thereof wherein W, Y, R₃, and R₅ are defined above.


18

The present invention also encompasses compounds of Formula V:



or the pharmaceutically acceptable non-toxic salts thereof wherein W, R₄, and R₅ are defined above.

The invention further encompasses compounds of Formula VI:



or the pharmaceutically acceptable non-toxic salts thereof where R₁₆ is hydrogen, benzyl or straight or branched chain lower alkyl having 1-6 carbon atoms;


R₁₇ is hydrogen or straight or branched chain lower alkoxy having 1-6 carbon atoms;

R₁₈ is hydrogen, straight or branched chain lower alkoxy having 1-6 carbon atoms, NR₁COR₂, COR₂, CO₂R₂ where R₁ and R₂ are as defined above, cyano, or halogen; and

R₁₉ is hydrogen or 42 halogen.

The invention additionally encompasses compounds of formula


VII:



VII

or the pharmaceutically acceptable non-toxic salts thereof
where R₁₇, R₁₈ and R₁₉ are defined above.


The invention further encompasses compounds of Formula VIII:



VIII

where R₁₇, R₁₈, and R₁₉ are defined above and R₂₀ is hydrogen or
straight or branched chain alkyl having 1-6 carbon atoms.


The invention further encompasses compounds of Formula IX:



IX


where R₁₇, R₁₈, and R₁₉ are defined above.

The invention further encompasses compounds of Formula X:



where R₁₆, R₁₇, R₁₈, and R₁₉ are defined above.


5 The invention further encompasses compounds of Formula XI:



where R₄, R₅, R₁₇, R₁₈, and R₁₉ are defined above.

10

The invention further encompasses compounds of Formula XII:



15

wherein W represents phenyl, 2 or 3-thienyl, 2, 3, or 4-pyridyl or 6-quinoliny, each of which may be mono or disubstituted with halogen, cyano, hydroxy, straight or branched chain lower alkyl having 1-6 carbon atoms, amino, mono or dialkylamino where each alkyl is

independently straight or branched chain lower alkyl having 1-6 carbon atoms, straight or branched chain lower alkoxy having 1-6 carbon atoms, or NR_1COR_2 , COR_2 , CONR_1R_2 or CO_2R_2 where R_1 and R_2 are the same or different and represent hydrogen or straight or branched chain lower alkyl having 1-6 carbon atoms

Non-toxic pharmaceutical salts include salts of acids such as hydrochloric, phosphoric, hydrobromic, sulfuric, sulfinic, formic, toluene sulfonic, hydroiodic, acetic and the like. Those skilled in the art will recognize a wide variety of non-toxic pharmaceutically acceptable addition salts.


Representative compounds of the present invention, which are encompassed by Formula I, include, but are not limited to the compounds in Figure 1 and their pharmaceutically acceptable salts. The present invention also encompasses the acylated prodrugs of the compounds of Formula I. Those skilled in the art will recognize various synthetic methodologies which may be employed to prepare non-toxic pharmaceutically acceptable addition salts and acylated prodrugs of the compounds encompassed by Formula I.

By lower alkyl in the present invention is meant straight or branched chain alkyl groups having 1-6 carbon atoms, such as, for example, methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, 2-pentyl, isopentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl, and 3-methylpentyl.

By lower alkoxy in the present invention is meant straight or branched chain alkoxy groups having 1-6 carbon atoms, such as, for example, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, sec-butoxy, tert-butoxy, pentoxy, 2-pentyl, isopentoxy, neopentoxy, hexoxy, 2-hexoxy, 3-hexoxy, and 3-methylpentoxy.

By halogen in the present invention is meant fluorine, bromine, chlorine, and iodine.

By N-alkylpiperazyl in the invention is meant radicals of the formula:



where R is a straight or branched chain lower alkyl as defined above.

The pharmaceutical utility of compounds of this invention are indicated by the following assay for GABA_A receptor binding activity.

Assays are carried out as described in Thomas and Tallman (J. Bio. Chem. 156: 9838-9842, J. Neurosci. 2: 433-440, 1983). Rat cortical tissue is

dissected and homogenized in 25 volumes²² (w/v) of 0.05 M Tris HCl buffer (pH 7.4 at 40°C). The tissue homogenate is centrifuged in the cold (4°C) at 20,000 x g for 20'. The supernatant is decanted and the pellet is rehomogenized in the same volume of buffer and again centrifuged at 20,000 x g. The supernatant
5 is decanted and the pellet is frozen at -20°C overnight. The pellet is then thawed and rehomogenized in 25 volume (original wt/vol) of buffer and the procedure is carried out twice. The pellet is finally resuspended in 50 volumes (w/vol) of 0.05 M Tris HCl buffer (pH 7.4 at 40°C).

Incubations contain 100 ml of tissue homogenate, 100 ml of
10 radioligand 0.5 nM (³H-RO15-1788 [³H-Flumazenil] specific activity 80 Ci/mmol), drug or blocker and buffer to a total volume of 500 ml. Incubations are carried for 30 min at 40°C then are rapidly filtered through GFB filters to separate free and bound ligand. Filters are washed twice with fresh 0.05 M Tris HCl buffer (pH 7.4 at 40°C) and counted in a liquid
15 scintillation counter. 1.0 mM diazepam is added to some tubes to determine nonspecific binding. Data are collected in triplicate determinations, averaged and % inhibition of total specific binding is calculated. Total Specific Binding = Total - Nonspecific. In some cases, the amounts of unlabeled drugs is varied and total displacement curves of binding are
20 carried out. Data are converted to a form for the calculation of IC₅₀ and Hill Coefficient (nH). Data for the compounds of this invention are listed in Table I.

23
TABLE I

| | <u>Compound Number</u> ¹ | <u>IC₅₀(μM)</u> |
|----|-------------------------------------|---|
| 5 | 1 | .004 |
| | 2 | .430 |
| | 6 | .001 |
| 10 | 8 | .025 |
| | 12 | .028 |
| 15 | 14 | .133 |
| | 17 | .470 |

¹Compound numbers relate to compounds shown in the figures.

20

The compounds of general formula I may be administered orally, topically, parenterally, by inhalation or spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used
25 herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition, there is provided a pharmaceutical formulation comprising a compound of general formula I and a pharmaceutically acceptable carrier. One or more compounds of general formula I may be present in association with one or more non-toxic
30 pharmaceutically acceptable carriers and/or diluents and/or adjuvants and if desired other active ingredients. The pharmaceutical compositions containing compounds of general formula I may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft
35 capsules, or syrups or elixirs.

Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents,
40 coloring agents and preserving agents in order to provide pharmaceutically

24
elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, 5 lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration 10 and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, 15 for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such 20 excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty 25 acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene 30 oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

35 Oily suspensions may be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and 40 flavoring agents may be added to provide palatable oral preparations. These

compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

Pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monoleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monoleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of general formula I may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the

26
rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.


Compounds of general formula I may be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration
5 used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anaesthetics, preservatives and buffering agents can be dissolved in the vehicle.

Dosage levels of the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-
10 indicated conditions (about 0.5 mg to about 7 g per patient per day). The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain between from about 1 mg to about 500 mg of an active
15 ingredient.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of
20 excretion, drug combination and the severity of the particular disease undergoing therapy.

An illustration of the preparation of compounds of the present invention is given in **Schemes I and II**.


α 7
 Scheme 1



where:

5 W, Z, and n are as defined above.

Scheme 11



5


W, R₃, Y, R₅, and R₆ are as defined above.

Those having skill in the art will recognize that the starting materials may be varied and additional steps employed to produce compounds encompassed by the present invention, as demonstrated by the following examples. In some cases protection of certain reactive functionalities may be necessary to achieve some of the above transformations. In general the need for such protecting groups will be apparent to those skilled in the art of organic synthesis as well as the conditions necessary to attach and remove such groups.

The invention is illustrated further by the following examples which are not to be construed as limiting the invention in scope or spirit to the specific procedures described in them.

29

Example 1




5

4-Oxo-4,5,6,7-tetrahydrobenzofuran-3-carboxylic acid was prepared according to a modified literature procedure ¹. To a stirring solution of potassium hydroxide (28.06 g, 0.5 mol) in methanol (100 mL) under Nitrogen at 0° C was added dropwise a solution of cyclohexanedione (56.07 g, 0.5 mol) in methanol (100 mL). The mixture was stirred at 0° C for 0.5 h, then a solution of ethyl bromopyruvate (66 mL, 0.525 mol) in methanol (100 mL) was added dropwise. After allowing the mixture to stir at ambient temperature for 17 h, a 50% aqueous sodium hydroxide solution (60 mL) was added dropwise and stirring continued an additional 7 h. After dilution with water, the solution was acidified and the methanol removed *in vacuo*. Ice water was added and the precipitate filtered and dried *in vacuo* to afford 4-Oxo-4,5,6,7-tetrahydrobenzofuran-3-carboxylic acid.

20

Example 2




To a stirring suspension of 4-Oxo-4,5,6,7-tetrahydrobenzofuran-3-carboxylic acid (28.2 g, 157 mmol) in ethanol (500 mL) under Nitrogen at ambient temperature was added acetyl chloride (56 mL, 783 mmol) dropwise. After stirring 1 h, the solution was then heated at reflux for 1 h. The solution was cooled and concentrated *in vacuo*. The residue was taken up into dichloromethane, washed with aqueous sodium bicarbonate, washed quickly with 1N sodium hydroxide, dried over magnesium sulfate, filtered, and concentrated *in vacuo* to afford Ethyl 4-oxo-4,5,6,7-

3a
tetrahydrobenzofuran-3-carboxylate as an oil. A mixture of this ester (24.46 g, 117 mmol) and ammonium acetate (15.85 g, 206 mmol) in dimethylformamide (225 mL) was heated at 100°C under Nitrogen for 1.25 h. The mixture was cooled, poured into ice water, and extracted two times with dichloromethane. The combined organic extracts were washed with water, dried over magnesium sulfate, filtered, concentrated *in vacuo*, and the residue triturated with ether to give Ethyl 4-oxo-4,5,6,7-tetrahydroindole-3-carboxylate. A mixture of this ester (11.31 g, 55 mmol) in 5N sodium hydroxide (200 mL) and ethanol (20 mL) was heated at reflux for 1h. After cooling in an ice bath, the mixture was acidified with concentrated hydrochloric acid, the precipitate filtered, rinsed with ice water, and dried *in vacuo* to afford 4-Oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxylic acid.

15

Example 3




Compound 1

20 A mixture of 4-Oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxylic acid (179 mg, 1 mmol), *p*-anisidine (616 mg, 5 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride [EDCI] (959 mg, 5 mmol) in 50% aqueous 1,4-dioxane (10 mL) was stirred at ambient temperature for 17 h. After concentrating *in vacuo*, the residue was taken up in 10% methanol in ethyl acetate, washed with 1.3M hydrochloric acid then with aqueous sodium bicarbonate, dried over magnesium sulfate, filtered, and concentrated *in vacuo*. Recrystallization from ethyl acetate afforded N-(4-Methoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 1); mp 219-220°C.

30


Example 4

2 /



- 5 A solution of 2-methyl-3-nitroanisole (9.96 g, 60 mmol) and tris(dimethylamino)methane (15.5 mL, 89 mmol) in dimethylformamide (30 mL) was heated at 115°C under Nitrogen for 3 h. After cooling to ambient temperature a solution of semicarbazide hydrochloride (6.98 g, 63 mmol) and concentrated hydrochloric acid (5.3 mL) in water (75 mL) was added
- 10 dropwise with vigorous stirring. The mixture was cooled further in an ice bath and the precipitate filtered, rinsed with ice water followed by, in sequence, cold 50% aqueous ethanol, cold ethanol, and ether, then dried to give semicarbazone. The semicarbazone was suspended with 10% Palladium on Carbon (3 g) in ethanol (120 mL) in a Parr shaker and placed under a
- 15 Hydrogen atmosphere (50 psi) for 16 h. The mixture was filtered through Celite and concentrated *in vacuo*. The residue was triturated with ice water, filtered and dried to give 4-methoxy-1H-indole.


~~30~~
Example 5



- 5 A solution of 2-nitro-6-benzyloxytoluene (13.0 g, 53 mmol) and tris(dimethylamino)methane (13.9 mL, 80 mmol) in dimethylformamide (30 mL) was heated at 115°C under Nitrogen for 3 h. Upon cooling to ambient temperature, a solution of semicarbazide hydrochloride (6.26 g, 56 mmol) and concentrated hydrochloric acid (4.8 mL) in water (70 mL) was added
- 10 dropwise with vigorous stirring. Ethanol (25 mL) was added and the heterogeneous mixture stirred for 2 h. After cooling in an ice bath, the precipitate was filtered, rinsed with, in sequence, ice water, cold 50% aqueous ethanol, cold ethanol, then ether and dried to give semicarbazone. A slurry of semicarbazone (17.64 g, 54 mmol), Raney Nickel (18 g of a 50%
- 15 aqueous slurry) in 1:1 tetrahydrofuran:methanol (145 mL) was heated to 55°C. Hydrazine monohydrate was added in four equal portions (2.7 mL each) at 0.5 h intervals. The mixture was cooled, then filtered through a small pad of silica gel using ether. The filtrate was dried over magnesium sulfate, filtered, concentrated *in vacuo*, and the residue purified by flash
- 20 chromatography to afford 4-benzyloxy-1H-indole as a low melting solid.

33

Example 6




5

To a solution of 4-methoxy-1H-indole (7.15 g, 49 mmol) and pyridine (19.7 mL, 243 mmol) in dichloromethane (50 mL) at 0°C under Nitrogen was added dropwise a solution of trichloroacetyl chloride (27.1 mL, 243 mmol) in dichloromethane (50 mL). After stirring at 0°C for an additional 1.5 h, the mixture was concentrated *in vacuo*. The residue was taken up in the minimal volume of methanol necessary and placed in a freezer overnight. The precipitate was filtered, rinsed well with methanol, and dried to yield 4-methoxy-3-trichloroacetyl-1H-indole. To a stirring solution of sodium methoxide (3.5 mL of a 25 weight % solution in methanol) in methanol (200 mL) at ambient temperature was added 4-methoxy-3-trichloroacetyl-1H-indole (9.07 g, 31.0 mmol) in portions over 0.75 h. After stirring an additional 0.75 h, the mixture was cooled in an ice bath, diluted with ice water, acidified with concentrated hydrochloric acid, and the methanol removed *in vacuo*. The resulting heterogeneous mixture was cooled in an ice bath, the precipitate filtered, rinsed with water, and dried to afford methyl 4-methoxy-1H-indole-3-carboxylic ester. A slurry of this ester (5.6 g 27 mmol) in 50% aqueous sodium hydroxide (50 mL) and methanol (50 mL) was stirred at ambient temperature for 19 h. The mixture was cooled in an ice bath, acidified with concentrated hydrochloric acid, the precipitate filtered, rinsed with ice water, and dried to afford 4-methoxy-1H-indole-3-carboxylic acid.

34

Example 7



(Compound 2)

5 N-(2-Fluoro-4-methoxyphenyl)-4-benzyloxy-1H-indole-3-carboxamide (1.34 g, 3.4 mmol), prepared using the method above, was slurried with 10% Palladium on Carbon (134 mg) in ethanol (35 mL) in a
10 Parr bottle and placed under a Hydrogen atmosphere (50 psi) for 5 h. Methanol (5 ml) was added and the mixture returned to the Hydrogen atmosphere for an additional 18 h. The solution was filtered through Celite, concentrated *in vacuo*, and the residue purified by flash chromatography to afford N-(2-fluoro-4-methoxyphenyl)-4-hydroxy-1H-indole-3-carboxamide
15 (Compound 2) as a beige solid; mp 259-261°C (d).

Example 8


20 The following compounds were prepared essentially according to the procedures described in Examples 1-6:

- (a) N-(4-Ethoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 3); mp 217-218°C.
- 25 (b) N-(4-Ethoxyphenyl)-4-methoxy-1H-indole-3-carboxamide (Compound 4); mp 259-261°C (d).
- (c) N-(3-Ethoxyphenyl)-4-methoxy-1H-indole-3-carboxamide (Compound 5).


30

- (d) N-(2-Fluoro-4-methoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 6).
- (e) N-(2-Fluoro-4-ethoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 7).
- (f) N-(4-Methoxyphenyl)-4-methoxy-1H-indole-3-carboxamide (Compound 8).
- (g) N-(2-Fluoro-4-ethoxyphenyl)-4-methoxy-1H-indole-3-carboxamide (Compound 9).
- (h) N-(4-Cyanophenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 10); mp 287-288°C.
- (i) N-(3-Methoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 11); mp 220-223°C.
- (j) N-(4-Acetamidophenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 12); mp 339-341°C.
- (k) N-(4-Chlorophenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 13).
- (l) N-(2-Fluoro-4-methoxyphenyl)-4-benzyloxy-1H-indole-3-carboxamide (Compound 14).
- (m) N-(4-acetylphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 15).
- (n) N-(2-Fluoro-5-methoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 16).
- (o) N-(6-quinoliny)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 17); mp 319-321°C.
- (p) N-(4-Carbomethoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 18); mp 261-262°C.


- (q) N-(4-Carboxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 19).
- (r) N-(4-Ethoxyphenyl)-4-hydroxy-1H-indole-3-carboxamide
5 (Compound 20).
- (s) N-(3-ethoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 21).
- (t) N-(2-fluoro-4-methoxyphenyl)-4-benzyloxy-1H-indole-3-carboxamide (Compound 22).
- (u) N-(4-isopropoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 23).
- 15 (v) N-(4-fluorophenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 24).
- (w) N-(4-pyridyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 25).
- 20 (x) N-(4-hydroxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 26).
- (y) N-(4-aminophenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 27).
- 25 (z) N-(3-pyridyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 28).
- 30 (aa) N-(4-(methoxymethyl)phenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 29).
- (ab) N-(3-fluoro-4-methoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 30).
- 35 (ac) (compound 31) (ad) (compound 32)




37



(ad) (compound 33)



(ae) (compound 34)



5

(af) N-(2-fluoro-4-isopropoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 35).

10 (ag) N-(4-fluorophenyl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 36).

(ah) N-(2-fluoro-4-methoxyphenyl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 37).

15 (ai) N-(4-ethoxyphenyl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 38).

(aj) N-(4-(methoxymethyl)phenyl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 39).

20

(ak) N-(4-methoxyphenyl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 40).

25 (al) N-(2-methoxy-5-pyridyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 41).

(am) N-(3,4-dihydroxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 42).

- 28
- (an) N-phenyl-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide
(Compound 43).
- (ao) N-(2-pyridyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-
5 carboxamide (Compound 44).
- (ap) N-(2-fluoro-4-hydroxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-
indole-3-carboxamide (Compound 45).
- 10 (aq) N-(2-fluoro-4-methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-
tetrahydro-1H-indole-3-carboxamide (Compound 46).
- (ar) N-(2-fluoro-4-ethoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-
tetrahydro-1H-indole-3-carboxamide (Compound 47).
- 15 (as) N-(4-ethoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-
indole-3-carboxamide (Compound 48).
- (at) N-(2-amino-3-pyridyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-
20 carboxamide (Compound 49).
- (au) N-(4-methylphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-
carboxamide (Compound 50).
- 25 (av) N-(3-methylphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-
carboxamide (Compound 51).
- (aw) N-(2-amino-4-methylphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-
indole-3-carboxamide (Compound 52).
- 30 (ax) N-(4-methylaminophenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-
3-carboxamide (Compound 53).
- (ay) N-phenyl-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-
35 carboxamide (Compound 54).
- (az) N-(3-hydroxy-4-methoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-
indole-3-carboxamide (Compound 55).

39

(ba) N-(2-fluorophenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 56).

(bb) N-(3-fluorophenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 57).

(bc) N-(4-(2-hydroxyethoxy)phenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 58).

(bd) N-(2-thienyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 59).


(be) N-(4-(2-methoxyethoxy)phenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 60).

15


(bf) N-(2-(4-hydroxyphenyl)ethyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 61).

(bg) N-(2-(4-ethoxyphenyl)ethyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 62).

(bh) (compound 63)




(bi) (compound 64)




25

(bj) (compound 65)


(bk) (compound 66)





(bl) (compound 67)



(bm) (compound 68)



(bn) (compound 69)



(bo) N-(4-fluoro-2-hydroxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-
10 indole-3-carboxamide (Compound 70).

5


The invention and the manner and process of making and using it, are now described in such full, clear, concise and exact terms as to enable any person skilled in the art to which it pertains, to make and use the same. It is to be understood that the foregoing describes preferred embodiments of the present invention and that modifications may be made therein without departing from the spirit or scope of the present invention as set forth in the claims. To particularly point out and distinctly claim the subject matter regarded as invention, the following claims conclude this specification.

10

WHAT IS CLAIMED IS:

42

1. A compound of the formula:



5

or the pharmaceutically acceptable non-toxic salts thereof wherein:

10 T is

halogen, hydrogen, hydroxyl, amino or straight or
branched chain lower alkoxy having 1-6 carbon atoms;

X is

15

hydrogen, hydroxyl or straight or branched chain lower
alkyl having 1-6 carbon atoms;

W is


20

phenyl, 2 or 3-thienyl, 2, 3, or 4-pyridyl or 6-
quinoliny, each of which may be mono or disubstituted with
halogen, cyano, hydroxy, straight or branched chain lower
alkyl having 1-6 carbon atoms, amino, mono or dialkylamino
where each alkyl is independently straight or branched chain
lower alkyl having 1-6 carbon atoms, straight or branched
chain lower alkoxy having 1-6 carbon atoms, or NR_1COR_2 , COR_2 ,
25 $CONR_1R_2$ or CO_2R_2 where R_1 and R_2 are the same or different
and represent hydrogen or straight or branched chain lower
alkyl having 1-6 carbon atoms; and

30



represents



wherein:

Y represents nitrogen or C- R_4 ;

Z represents N- R_7 or a carbon atom substituted with R_8 and R_9 , i.e., $C(R_8)(R_9)$;

n is 1, 2, 3, or 4;

R_3 is

hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2, 3 or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms;

R_4 is

halogen or trifluoromethyl; or

- OR_{10} , - COR_{10} , - CO_2R_{10} , - $OCOR_{10}$, or - R_{10} , where R_{10} is hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2, 3 or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; or

- $CONR_{11}R_{12}$ or $-(CH_2)_mNR_{11}R_{12}$, where m is 0, 1, or 2; R_{11} represents hydrogen, straight or branched chain lower alkyl having 1-6 carbon atoms; and R_{12} is hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2, 3 or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; or $NR_{11}R_{12}$ forms a

heterocyclic group which is morpholyl, piperidyl, pyrrolidyl, or N-alkyl piperazyl;

R₅ and R₆ are the same or different and represent

5 hydrogen, halogen, straight or branched chain lower alkyl having 1-6 carbon atoms, or straight or branched chain lower alkoxy having 1-6 carbon atoms;

R₇ is

10 hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2,3, or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms;

15

R₈ is

hydrogen or straight or branched chain lower alkyl having 1-6 carbon atoms; and

20

R₉ is

-COR₁₃, -CO₂R₁₃ or -R₁₃, where R₁₃ is hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2, 3, or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; or


25

-CONR₁₄R₁₅ or -(CH₂)_kNR₁₄R₁₅, where k is 0, 1, or 2; R₁₄ represents hydrogen, straight or branched chain lower alkyl having 1-6 carbon atoms; and R₁₅ is hydrogen, phenyl, 2, 3, or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2, 3 or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; or NR₁₄R₁₅ forms a heterocyclic group which is morpholyl, piperidyl, pyrrolidyl, or N-alkyl piperazyl.

30

35

40 2. A compound of the formula:



or the pharmaceutically acceptable non-toxic salts thereof wherein:

5

W is

phenyl, 2 or 3-thienyl, 2, 3, or 4-pyridyl or 6-quinoliny, each of which may be mono or disubstituted with
 10 halogen, cyano, hydroxy, straight or branched chain lower alkyl having 1-6 carbon atoms, amino, mono or dialkylamino where each alkyl is independently straight or branched chain lower alkyl having 1-6 carbon atoms, straight or branched chain lower alkoxy having 1-6 carbon atoms, or NR_1COR_2 , COR_2 ,
 15 CONR_1R_2 or CO_2R_2 where R_1 and R_2 are the same or different and represent hydrogen or straight or branched chain lower alkyl having 1-6 carbon atoms; and

R_8 is

20 hydrogen or straight or branched chain lower alkyl having 1-6 carbon atoms; and

R_9 is


25 $-\text{COR}_{13}$, $-\text{CO}_2\text{R}_{13}$ or $-\text{R}_{13}$, where R_{13} is hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2, 3, or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; or

30 $-\text{CONR}_{14}\text{R}_{15}$ or $-(\text{CH}_2)_k\text{NR}_{14}\text{R}_{15}$, where k is 0, 1, or 2; R_{14} represents hydrogen, straight or branched chain lower alkyl having 1-6 carbon atoms; and R_{15} is hydrogen, phenyl, 2, 3, or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2, 3 or 4-pyridylalkyl where each alkyl is straight or branched chain
 35 lower alkyl having 1-6 carbon atoms; or $\text{NR}_{14}\text{R}_{15}$ forms a

heterocyclic group which is morpholyl, piperidyl, pyrrolidyl, or N-alkyl piperazyl.

3. A compound of the formula:

5



or the pharmaceutically acceptable non-toxic salts thereof wherein:

10

W is

phenyl, 2 or 3-thienyl, 2, 3, or 4-pyridyl or 6-quinolinyl, each of which may be mono or disubstituted with halogen, cyano, hydroxy, straight or branched chain lower alkyl having 1-6 carbon atoms, amino, mono or dialkylamino where each alkyl is independently straight or branched chain lower alkyl having 1-6 carbon atoms, straight or branched chain lower alkoxy having 1-6 carbon atoms, or NR_1COR_2 , COR_2 , CONR_1R_2 or CO_2R_2 where R_1 and R_2 are the same or different and represent hydrogen or straight or branched chain lower alkyl having 1-6 carbon atoms; and

15

20

R_3 is

hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2, 3 or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; and


25

30

R_5 is

hydrogen, halogen, straight or branched chain lower alkyl having 1-6 carbon atoms, or straight or branched chain lower alkoxy having 1-6 carbon atoms.

35 4. A compound of the formula:



5 and the pharmaceutically acceptable non-toxic salts thereof wherein:

W is

10 phenyl, 2 or 3-thienyl, 2, 3, or 4-pyridyl or 6-quinoliny, each of which may be mono or disubstituted with halogen, cyano, hydroxy, straight or branched chain lower alkyl having 1-6 carbon atoms, amino, mono or dialkylamino where each alkyl is independently straight or branched chain lower alkyl having 1-6 carbon atoms, straight or branched chain lower alkoxy having 1-6 carbon atoms, or NR₁COR₂, COR₂, 15 CONR₁R₂ or CO₂R₂ where R₁ and R₂ are the same or different and represent hydrogen or straight or branched chain lower alkyl having 1-6 carbon atoms; and


R₃ is

20 hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2, 3 or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; and

25 R₅ is

hydrogen, halogen, straight or branched chain lower alkyl having 1-6 carbon atoms, or straight or branched chain lower alkoxy having 1-6 carbon atoms.

30 5. A compound of the formula:



and the pharmaceutically acceptable non-toxic salts thereof wherein:

5 W is

phenyl, 2 or 3-thienyl, 2, 3, or 4-pyridyl or 6-quinoliny, each of which may be mono or disubstituted with halogen, cyano, hydroxy, straight or branched chain lower alkyl having 1-6 carbon atoms, amino, mono or dialkylamino
10 where each alkyl is independently straight or branched chain lower alkyl having 1-6 carbon atoms, straight or branched chain lower alkoxy having 1-6 carbon atoms, or NR₁COR₂, COR₂, CONR₁R₂ or CO₂R₂ where R₁ and R₂ are the same or different and represent hydrogen or straight or branched chain lower
15 alkyl having 1-6 carbon atoms; and

Y is

nitrogen or C-R₄;

R₄ is

20 halogen or trifluoromethyl; or

-OR₁₀, -COR₁₀, -CO₂R₁₀, -OCOR₁₀, or -R₁₀, where R₁₀ is hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or
25 2, 3 or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; or

-CONR₁₁R₁₂ or -(CH₂)_mNR₁₁R₁₂, where m is 0, 1, or 2; R₁₁ represents hydrogen, straight or branched chain lower alkyl having 1-6 carbon atoms; and R₁₂ is hydrogen, phenyl,
30 2, 3 or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2, 3 or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; or NR₁₁R₁₂ forms a

49


heterocyclic group which is morpholyl, piperidyl, pyrrolidyl, or N-alkyl piperazyl; and

R₅ is

- 5 hydrogen, halogen, straight or branched chain lower alkyl having 1-6 carbon atoms, or straight or branched chain lower alkoxy having 1-6 carbon atoms.

6. A compound of the formula:

10



and the pharmaceutically acceptable non-toxic salts thereof wherein:

15 W is

- phenyl, 2 or 3-thienyl, 2, 3, or 4-pyridyl or 6-quinoliny, each of which may be mono or disubstituted with halogen, cyano, hydroxy, straight or branched chain lower alkyl having 1-6 carbon atoms, amino, mono or dialkylamino where each alkyl is independently straight or branched chain lower alkyl having 1-6 carbon atoms, straight or branched chain lower alkoxy having 1-6 carbon atoms, or NR₁COR₂, COR₂, CONR₁R₂ or CO₂R₂ where R₁ and R₂ are the same or different and represent hydrogen or straight or branched chain lower alkyl having 1-6 carbon atoms; and
- 20
- 25

R₄ is

halogen or trifluoromethyl; or

- 30 -OR₁₀, -COR₁₀, -CO₂R₁₀, -OCOR₁₀, or -R₁₀, where R₁₀ is hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2, 3 or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; or
- 35

$\text{-CONR}_{11}\text{R}_{12}$ or $\text{-(CH}_2\text{)}_m\text{NR}_{11}\text{R}_{12}$, where m is 0, 1, or 2; R_{11} represents hydrogen, straight or branched chain lower alkyl having 1-6 carbon atoms; and R_{12} is hydrogen, phenyl, 2, 3 or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2, 3 or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; or $\text{NR}_{11}\text{R}_{12}$ forms a heterocyclic group which is morpholyl, piperidyl, pyrrolidyl, or N-alkyl piperazyl; and

R_5 is

hydrogen, halogen, straight or branched chain lower alkyl having 1-6 carbon atoms, or straight or branched chain lower alkoxy having 1-6 carbon atoms.

15

7. A compound according to claim 1, which is N-(4-Methoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

20 8. A compound according to claim 1, which is N-(2-fluoro-4-methoxyphenyl)-4-hydroxy-1H-indole-3-carboxamide.

9. A compound according to claim 1, which is N-(4-Ethoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

25 10. A compound according to claim 1, which is N-(4-Ethoxyphenyl)-4-methoxy-1H-indole-3-carboxamide.

11. A compound according to claim 1, which is N-(3-Ethoxyphenyl)-4-methoxy-1H-indole-3-carboxamide.

30


12. A compound according to claim 1, which is N-(2-Fluoro-4-methoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

35 13. A compound according to claim 1, which is N-(2-Fluoro-4-ethoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.


14. A compound according to claim 1, which is N-(4-Methoxyphenyl)-4-methoxy-1H-indole-3-carboxamide.

15. A compound according to claim 1, which is N-(2-Fluoro-4-ethoxyphenyl)-4-methoxy-1H-indole-3-carboxamide.
16. A compound according to claim 1, which is N-(4-Cyanophenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
17. A compound according to claim 1, which is N-(3-Methoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
18. A compound according to claim 1, which is N-(4-Acetamidophenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
19. A compound according to claim 1, which is N-(4-Chlorophenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
20. A compound according to claim 1, which is N-(2-Fluoro-4-methoxyphenyl)-4-benzyloxy-1H-indole-3-carboxamide.
21. A compound according to claim 1, which is N-(4-acetylphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
22. A compound according to claim 1, which is N-(2-Fluoro-5-methoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
23. A compound according to claim 1, which is N-(6-quinoliny)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
24. A compound according to claim 1, which is N-(4-Carbomethoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
25. A compound according to claim 1, which is N-(4-Carboxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
26. A compound according to claim 1, which is N-(4-Ethoxyphenyl)-4-hydroxy-1H-indole-3-carboxamide.
27. A compound according to claim 1, which is N-(3-ethoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.


28. A compound according to claim 1, which is N-(2-fluoro-4-methoxyphenyl)-4-benzyloxy-1H-indole-3-carboxamide.
29. A compound according to claim 1, which is N-(4-isopropoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
30. A compound according to claim 1, which is N-(4-fluorophenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
31. A compound according to claim 1, which is N-(4-pyridyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
32. A compound according to claim 1, which is N-(4-hydroxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
33. A compound according to claim 1, which is N-(4-aminophenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
34. A compound according to claim 1, which is N-(3-pyridyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
35. A compound according to claim 1, which is N-(4-(methoxymethyl)phenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
36. A compound according to claim 1, which is N-(3-fluoro-4-methoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
37. A compound according to claim 1, which is



38. A compound according to claim 1, which is




39. A compound according to claim 1, which is



5

40. A compound according to claim 1, which is



41. A compound according to claim 1, which is N-(2-fluoro-4-isopropoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.


42. A compound according to claim 1, which is N-(4-fluorophenyl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

43. A compound according to claim 1, which is N-(2-fluoro-4-methoxyphenyl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.


44. A compound according to claim 1, which is N-(4-ethoxyphenyl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

- 34
45. A compound according to claim 1, which is N-(4-(methoxymethyl)phenyl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 5 46. A compound according to claim 1, which is N-(4-methoxyphenyl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
47. A compound according to claim 1, which is N-(2-methoxy-5-pyridyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 10 48. A compound according to claim 1, which is N-(3,4-dihydroxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
49. A compound according to claim 1, which is N-phenyl-4-oxo-4,5,6,7-
15 tetrahydro-1H-indole-3-carboxamide.
50. A compound according to claim 1, which is N-(2-pyridyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 20 51. A compound according to claim 1, which is N-(2-fluoro-4-hydroxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
52. A compound according to claim 1, which is N-(2-fluoro-4-methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-
25 carboxamide.
53. A compound according to claim 1, which is N-(2-fluoro-4-ethoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-
carboxamide.
- 30 54. A compound according to claim 1, which is N-(4-ethoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
55. A compound according to claim 1, which is N-(2-amino-3-pyridyl)-4-
35 oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
56. A compound according to claim 1, which is N-(4-methylphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

57. A compound according to claim 1, which is N-(3-methylphenyl)-4-
oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
58. A compound according to claim 1, which is N-(2-amino-4-
5 methylphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
59. A compound according to claim 1, which is N-(4-
methylaminophenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 10 60. A compound according to claim 1, which is N-phenyl-4-oxo-6,6-
dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
61. A compound according to claim 1, which is N-(3-hydroxy-4-
methoxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 15 62. A compound according to claim 1, which is N-(2-fluorophenyl)-4-oxo-
4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
63. A compound according to claim 1, which is N-(3-fluorophenyl)-4-oxo-
20 6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
64. A compound according to claim 1, which is N-(4-(2-
hydroxyethoxy)phenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 25 65. A compound according to claim 1, which is N-(2-thienyl)-4-oxo-
4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
66. A compound according to claim 1, which is N-(4-(2-
methoxyethoxy)phenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 30 67. A compound according to claim 1, which is N-(2-(4-
hydroxyphenyl)ethyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
68. A compound according to claim 1, which is N-(2-(4-
35 ethoxyphenyl)ethyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
69. A compound according to claim 1, which is




70. A compound according to claim 1, which is




5


71. A compound according to claim 1, which is



72. A compound according to claim 1, which is




73. A compound according to claim 1, which is




5

74. A compound according to claim 1, which is




75. A compound according to claim 1, which is




76. A compound according to claim 1, which is N-(4-fluoro-2-hydroxyphenyl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

Fig. 1




Compound 1

Fig. 2




Compound 2

Fig. 3



Compound 6


Fig. 4



Compound 8


2 / 5

Fig. 5




Compound 12

Fig. 6




Compound 14

Fig. 7




Compound 17

Fig. 8




Compound 3

Fig. 9




Compound 4

Fig. 10




Compound 5

Fig. 11




Compound 7

Fig. 12




Compound 9

Fig. 13




Compound 10

Fig. 14




Compound 11

Fig. 15



Compound 13

Fig. 16



Compound 15

5 / 5

Fig. 17






Fig. 18



Compound 16


Compound 18

Fig. 19



Compound 19

Fig. 20



Compound 20

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D209/42 C07D209/52 C07D401/12 C07D413/04 C07D405/12
C07D409/12 C07D471/04 A61K31/44 A61K31/40

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| A | US, A, 5 243 049 (K. SHAW ET AL.) 7 September 1993 see the whole document ----- | 1-76 |

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another claim or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"A" document member of the same patent family

Date of the actual completion of the international search

27 December 1994

Date of mailing of the international search report

- 4. 01. 95

Name and mailing address of the ISA

European Patent Office, P.B. 3818 Patentaan 2
NL - 2200 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tlx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Bosma, P

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 93/12300

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| US-A-5243049 | 07-09-93 | WO-A- 9426741 | 24-11-94 |